

## Genetic Diversity in Black Howler Monkeys (*Alouatta pigra*) From Belize

ROXIE A. JAMES,<sup>1,4\*</sup> PAUL L. LEBERG,<sup>2</sup> JOSEPH M. QUATTRO,<sup>3</sup>  
AND ROBERT C. VRIJENHOEK<sup>4</sup>

<sup>1</sup>Department of Biology, Kean College of New Jersey, Union,  
New Jersey 07083-0411

<sup>2</sup>Department of Biology, University of Southwestern Louisiana,  
Lafayette, Louisiana 70504

<sup>3</sup>Department of Biology, University of South Carolina, Columbia,  
South Carolina 29208

<sup>4</sup>Center for Theoretical and Applied Genetics Rutgers University,  
New Brunswick, NJ 08903-0231

**KEY WORDS** allozymes; restriction fragments; threatened species; bottlenecks; mitochondrial DNA; mtDNA; heterozygosity; population structure

**ABSTRACT** To assess the level of genetic variation in a threatened black howler monkey (*Alouatta pigra*) population, we examined 36 allozyme loci and restriction fragment profiles of mitochondrial DNA (mtDNA). Mean heterozygosity at allozyme loci was only 0.021 and 5.6 percent of the loci were polymorphic. Analyses of mtDNA also revealed low genetic diversity compared with other primates. F-statistics revealed no significant genetic heterogeneity among troops within the Bermudian Landing preserve, but did indicate a deficiency of heterozygotes at one of the two loci. We explore several explanations for this result, which is unexpected in a socially structured primate. Low genetic diversity in this population may reflect its history of demographic bottlenecks. *Am. J. Phys. Anthropol.* 102:329-336, 1997.

© 1997 Wiley-Liss, Inc.

Belizean black howler monkeys (*Alouatta pigra*, Lawrence, 1933) have undergone four population crashes in recent times. Devastating hurricanes swept through Bermudian Landing, Belize in 1931, 1954, and 1978 (Hartshorn, 1984; Bolin, 1981), and a yellow fever epidemic decimated Central American howlers in 1971 (Baldwin, 1976; Hartshorn, 1984). Each of these events reduced local populations to very low levels, affecting both the sizes and the behavioral dynamics of remnant troops (Baldwin, 1976; Bolin, 1981). Although remnant populations have recovered, troops remain small compared to other howler species such as Costa Rican golden mantled howlers (*Alouatta palliata*, Gray, 1849) in which troop size averages about 20 (Glander, 1980). Troop composition in black howlers varies but generally consists of one

dominant adult male, one or two adult females, plus subadults and juveniles (mean troop size = 4.6, S.D. = 1.7).

When a population experiences a demographic bottleneck, genetic diversity is expected to decline depending upon the bottleneck population size and the subsequent rate of population growth (Chakraborty and Nei, 1977; Maruyama and Fuerst, 1985a,b;

Contract grant sponsor: New Jersey Agricultural Experiment Station; Contract grant number D-67175-x-94; Contract grant sponsor: NSF; Contract grant number INT84-16427; Contract grant number BSR86-00661; Contract grant number BSR88-05361; Contract grant number DEB-9123943; Contract grant number 9316246; Contract grant sponsor: National Geographic Society; Contract grant number 4773-92; Contract grant sponsor: AAUW.

\*Correspondence to: Roxie A. James, Department of Biology, Kean College of New Jersey, Union, NJ 07083.

Received 9 May 1994; revised 23 April 1996; accepted 27 November 1996.

Nei et al., 1975; Leberg, 1992). A demographic bottleneck can result from either a reduction in the size of a population to a few individuals or from the establishment of a population by a small number of founders. Given the recent history of bottlenecks in the Bermudian Landing black howlers, we would expect low levels of heterozygosity and allelic diversity compared with larger more stable populations. Barring immigration, it would take a very long time to restore variability in a troop through accumulated mutation (Sirrkomaa, 1983).

Social structure in Belizean howler populations might affect levels of genetic diversity in and among troops. Long-term behavioral studies identified a number of troops, each of which constitutes a stable assemblage of individuals with respect to home range (Horwich, 1983a,b; Horwich and Lyon, 1990). Social structure which imposes deviations from random mating and reduces dispersal among local breeding units can have significant effects on the distribution and amount of genetic variation in mammalian populations (Bush et al., 1977; Chesser, 1991a,b; Melnick et al., 1984; Smouse and Neel, 1981). For example, social structure can result in genetic differences among troops, as is seen in the red howler (*Alouatta seniculus*; Pope, 1992). The effects of population bottlenecks on genetic diversity are also dependent on social structure. In a demic model with limited gene flow among subpopulations, demographic bottlenecks decrease genetic diversity within the subpopulations but increase genetic differences among them. However, in populations comprised of social groups of philopatric females and dispersing males, demographic bottlenecks decrease both within-group diversity and genetic differences among groups (Chesser, 1991a).

The primary purpose of this paper is to examine genetic diversity and population structure in Belizean black howlers (*Alouatta pigra*) using allozyme loci and mitochondrial DNA (mtDNA). Our assessment of genetic diversity within and among social groups should provide useful information to conservation biologists attempting to preserve this species, as well as insight into the effects of population bottlenecks on the evo-

lution of socially structured populations of primates.

## MATERIALS AND METHODS

Howler troops were studied in the Community Baboon Sanctuary (CBS), Bermudian Landing, Belize. These troops had been defined previously on the basis of behavioral studies (Horwich, 1983a,b). The CBS is an area of second-growth forest; the sanctuary is not isolated from nearby habitat supporting howler troops.

Tissue samples were collected from all black howlers occupying the Community Baboon Sanctuary, during January–February 1990. We used standard dart gun techniques (Cohen and Bree, 1978; Glander, 1991; Scott et al., 1976) to anesthetize juveniles and adults within their habitats. Darts were loaded with 25 mg of Telazol (Tiletamine hydrochloride and zolazepam hydrochloride, A.H. Robbins Co., Richmond, VA) per kg of body weight. Animals falling from trees were caught in a nylon mesh net. Once on the ground, we took a blood sample from the femoral vein of all individuals >6 months of age. Blood was held in a Vacutainer (Becton Dickinson, Co., Rutherford, NJ) syringe with the anticoagulant EDTA. A small notch was cut in the ear lobe to mark each individual, regardless of age. We placed the ear tissue in a sterile 1.5 ml microcentrifuge tube and froze it immediately on dry ice. The anesthetized monkeys were placed in burlap bags in the shade until they recovered from the anesthesia.

The blood samples were kept on water ice packs for up to 6 hours in the field. Each evening at the field station, the fresh blood samples were centrifuged at 1,500 rpm for 20 minutes to separate plasma, erythrocytes, and leukocytes. The erythrocyte subsample was washed three times in 0.9% saline and suspended in an equal volume of mannitol-glycerol-saline solution to prevent lysis (Melnick et al., 1984). All samples were packed on dry ice and held at  $-20^{\circ}\text{C}$  for up to 10 days prior to transport to Rutgers University.

In the laboratory, samples of red cells were thawed and lysed with an equal volume of ice-cold grinding buffer, Tris-EDTA-MnCl<sub>2</sub>, pH 7.0. Plasma cells were lysed by

centrifugation (Harris and Hopkinson, 1978; Melnick et al., 1984). Extracts were loaded into preformed wells of 12% starch gels in the appropriate buffer (Table 1). Standard procedure for starch gel electrophoresis follows Harris and Hopkinson (1978) and Murphy et al. (1990). The designations used for gene loci follow those of Harris and Hopkinson (1978).

Allelic frequencies and tests for deviations from Hardy-Weinberg expectations were calculated using the computer program BIOSYS-1 (Swofford and Selander, 1981). Estimation of population structure were made using Wright's F-statistics (1965) with modifications following Nei (1977). One solitary male was not used in the estimation of the F-statistics because it could not be assigned to a troop. Because almost the entire population (all members >6 months of age) was sampled in terms of both troops and individuals, we did not use the approach of Weir and Cockerham (1984), which corrects for error due to incomplete sampling of demes and individuals. Following the suggestion of Chesson (1991a), we attempted to make separate estimates of the F-statistics for the adults, subadults, and juveniles; unfortunately, the number of individuals in the troops was too small to conduct this analysis.

We tested the significance of  $F_{IT}$ , the correlation of uniting gametes within individuals relative to that of the total population,  $F_{IS}$ , the average correlation of uniting gametes within individuals relative to that of a subdivision (troop, in our case), and  $F_{ST}$ , the correlation of uniting gametes within troops relative to that of the total population, using a randomization approach similar to that used by Pope (1992). The only deviation from her approach is that alleles, not genotypes, found in the 39 individual troop members, were randomly reassigned to individuals in the 10 troops. By randomly reassigning alleles to individuals and troops, rather than reassigning genotypes to troops, we could evaluate the significance of  $F_{IT}$ , as well as the other two components of genetic variance. This method was used to evaluate the significance of F-statistics for single loci, and mean values based on multiple loci; the results which these hypotheses test were not

TABLE 1. Proteins encoded by 36 putative gene loci in *Belizian Alouatta pigra*. Enzyme systems are listed by tissue preparations that provided best resolution

Protein	EC number	Code	Buffer <sup>1</sup>
<b>Plasma</b>			
Adenosine deaminase II	3.3.4.4	<i>Ada-I</i>	1
Aspartate aminotransferase	2.6.1.1	<i>Aat-I</i>	2
Carbonic Anhydrase	4.2.1.1	<i>Ca</i>	4
Creatine Kinase II	2.7.4.3	<i>Ck-II</i>	2
Esterase C	3.1.1.-	<i>Es-C</i>	4
Esterase D	3.1.1.-	<i>Es-D</i>	4
Glutamate Dehydrogenase	1.1.1.47	<i>Gdh</i>	2
Peptidase A	3.4.11	<i>Pepd-A</i>	4
Peptidase C	3.4.11	<i>Pepd-C</i>	4
Peptidase E	3.4.11	<i>Pepd-E</i>	4
Phosphoglucosaminidase	2.7.5.1	<i>Pgm-II</i>	4
<b>Erythrocyte (1:10 dilution)</b>			
Adenylate kinase	2.7.4.3	<i>Adk</i>	2
Catalase	1.11.1.6	<i>Cat</i>	2
Creatine kinase I	2.7.4.3	<i>Ck-I</i>	2
Malate dehydrogenase I	1.1.1.37	<i>Mdh</i>	4
Malic enzyme	1.1.1.37	<i>Me</i>	2
Phosphoglucose isomerase I	5.3.19	<i>Pgi-I</i>	6
<b>Erythrocyte (undiluted)</b>			
Adenosine deaminase I	3.5.4.4	<i>Ada-I</i>	1
Albumin		<i>Alb</i>	1
Esterase A	3.1.1.-	<i>Es-A</i>	4
Esterase B	3.1.1.-	<i>Es-B</i>	4
General protein I		<i>Gp-I</i>	1
General protein II		<i>Gp-II</i>	1
Glucose-2-dehydrogenase	1.1.1.8	<i>G2dh</i>	2
Glucose-6-phosphate dehydrogenase	1.1.1.49	<i>G6pdh</i>	3
α-Glycerol phosphate dehydrogenase	1.1.1.43	<i>αGpd</i>	3
Hemoglobin		<i>Hb</i>	1
Lactate dehydrogenase A	1.1.1.27	<i>Ldh-A</i>	3
Lactate dehydrogenase B	1.1.1.27	<i>Ldh-B</i>	3
Mannose phosphate isomerase	5.3.1.8	<i>Mpi</i>	2
NADH-diaphorase	1.8.1.4	<i>Dia</i>	5
Peptidase B	3.4.11	<i>Pepd-B</i>	4
Peptidase D	3.4.13.9	<i>Pepd-D</i>	4
Phosphoglucosaminidase I	2.7.5.1	<i>Pgm-I</i>	4
Purine nucleoside phosphorylase	2.4.2.1	<i>Pnp</i>	7
Superoxide dismutase	1.15.1.1	<i>Sod</i>	3

<sup>1</sup> Buffers: 1, Tris-Boric Acid-EDTA (TBE 8.7); 2, Tris-Citric Acid (TC 6.8); 3, Tris-maleic acid-EDTA (TME 7.4); 4, Amino-propylmorpholine (AP 6.0); 5, Tris-citric acid (TC 6.0); 6, Tris versene borate (TVB 8.0); 7, Lithium hydroxide-boric acid (LHB 7.2). Buffers are described in Harris and Hopkinson (1978) and Murphy et al. (1990).

TABLE 2. Troop size (*N*), sample size (*n*), frequencies of common alleles and numbers of observed and expected heterozygotes (*H<sub>o</sub>* and *H<sub>e</sub>*) at two diallelic loci (*Pgi-1*, *Pnp*) in Bermudian Land troops of *Alouatta pigra*

Troop	N	n	<i>Pgi-I</i>			<i>Pnp</i>		
			Freq.	<i>H<sub>o</sub></i>	<i>H<sub>e</sub></i>	Freq.	<i>H<sub>o</sub></i>	<i>H<sub>e</sub></i>
Fig tree	4	3	0.67	0	1.3	0.50	1	1.5
Wade pasture	5	3	0.67	0	1.3	0.50	1	1.5
Cashew	4	3	0.33	0	1.3	0.83	1	0.8
Della	7	7	0.64	3	3.2	0.57	2	3.4
Y	4	4	0.75	2	1.5	0.63	3	1.9
Northern A	5	3	0.37	1	1.4	0.37	2	1.4
A	4	4	0.63	1	1.9	0.25	2	1.5
Peninsula	6	6	0.83	2	1.7	0.75	3	2.3
Baptist	3	3	0.33	2	1.3	0.50	3	1.5
School	4	3	0.50	1	1.5	0.83	1	0.8
Mean <sup>1</sup>	4.6	3.9	0.60	1.5	1.9	0.58	2.0	1.9

<sup>1</sup> Means weighted by sample size of troops.

different from those obtained from more traditional tests of the significance of F-statistics (i.e., Workman and Niswander, 1970). Five thousand randomizations of the data were used to assess statistical significance at the 0.05 level.

Mitochondrial DNA variation was revealed by restriction analysis using 13 endonucleases (AVAI, AVAIL, BAMHI, BGLII, ECORI, ECORV, HINCII, HINDIII, PVUII, PSTI, SCAI, XBAI, XHOI). Ear tissue was used in this analysis so mtDNA genotypes were available for all 47 individuals inhabiting the CBS, not just those over >6 months of ages used in the allozyme survey. The specific conditions of DNA extraction, restriction enzyme digestion, electrophoresis, and DNA transfer and visualization are described by Lansman et al. (1981) and Southern (1975). A heterologous mtDNA probe was isolated from the whole liver of a red howler monkey. The probe DNA was labeled by random primed incorporation of dioxigenin-labeled deoxyuridine-triphosphate (Genius, BMB). The number of nucleotide substitutions, and the divergence of between DNA sequences, was estimated for haplotypes with different patterns of restriction fragments (Nei, 1987; Nei and Li, 1979).

## RESULTS

### Allozyme variation

Based on comparison with human isozyme patterns (Harris and Hopkinson, 1978), we inferred that the 24 proteins surveyed (Table 1) are encoded by 36 putative gene loci. Only two loci (*Pgi-I* and *Pnp*) were

TABLE 3. F-statistics at polymorphic gene loci in Bermudian Landing troops of *Alouatta pigra*

Locus	<i>F<sub>IT</sub></i>	<i>F<sub>ST</sub></i>	<i>F<sub>IS</sub></i>
<i>Pgi-I</i>	0.436*	0.116	0.362*
<i>Pnp</i>	0.056	0.135	-0.091
Mean	0.246	0.125	0.138

\* *P* < 0.05.

polymorphic in our sample of *A. pigra* (Table 2). Polymorphism occurred in all ten troops, so the percentage of polymorphic loci (*P*) was 5.6%. Mean heterozygosity (*H*) varied from troop to troop, but averaged 0.021.

There was no significant genetic differentiation among the ten troops; estimated *F<sub>ST</sub>* for the two polymorphic loci were not significantly different from zero (Table 3). Likewise, average *F<sub>IT</sub>* and *F<sub>IS</sub>* were not different from zero; however, both *F<sub>IS</sub>* and *F<sub>IT</sub>* were larger than zero for *Pgi-I* locus (Table 3). The large *F<sub>IS</sub>* and *F<sub>IT</sub>* values for this locus were due to excess homozygosity within troops (Table 2). This high homozygosity was not due to a scoring bias against *Pgi-I* heterozygotes, as every individual was examined electrophoretically at least twice, and heterozygotes were unequivocal. The presence of an enzymatically silent *Pgi-I* allele could be responsible for the high homozygosity observed at this locus, but we observed no apparent homozygotes for a silent allele. All individuals produced a *Pgi-I* phenotype on the gels.

### mtDNA diversity

Restriction enzymes with four base-pair recognition sequences generated 25 to 30

TABLE 4. A partial survey of studies reporting two indices of genetic variation, the percentage of polymorphic loci ( $P\%$ ) and average multiple-locus heterozygosity ( $H$ ) in primates

Population	Sample size	Number troops	Number loci	$P\%$	$H$	Ref. <sup>1</sup>
<i>Alouatta pigra</i> , Belize	41	10	36	5.6	0.021	1
<i>A. seniculus</i> , Venezuela	137	18	29	32.7	0.099	2
<i>Macaca senica</i> , Sri Lanka	131	16	32	34.3	0.078	3
<i>M. fuscata</i> , Japan	1642	33	32	9.2	0.013	4
<i>M. fuscata</i> , Japan	35	01	33	18.2	0.044	5
<i>M. fuscata yakui</i> , Yaku Island	29	15	29	00.0	0.000	6
<i>M. fascicularis</i> , Indonesia	456	29	33	12.2	0.038	7
<i>M. mulatta</i> , Pakistan	189	05	35	14.3	0.055	9
<i>Theropithecus gelada</i> , Ethiopia	213	04	34	7.4	0.013	10
<i>Papio hemadryas</i> , Ethiopia	414	05	34	31.4	0.042	11
<i>P. anubis</i> , Ethiopia	40	02	34	31.4	0.032	11
<i>Cercopithecus aethiops</i> , Ethiopia	124	07	23	17.4	0.056	12
<i>Homo sapiens</i> , Caucasian			104	23.0	0.063	13

<sup>1</sup> References: 1, present study; 2, Pope (1992); 3, Shotake and Santiapillai (1982); 4, Nozawa et al. (1982); 5, Hayasaka (1987); 6, Nozawa et al. (1977); 7, Kawamoto et al. (1984); 8, Ishimoto (1973); 9, Melnick et al. (1984); Melnick (1987); 10, Shotake and Nozawa (1984); 11, Shotake (1981); 12, Turner (1981); 13, Harris and Hopkinson (1978).

fragments per individual and produced fragment profiles that were uninterpretable. We limited our analysis to ten enzymes that recognize six base-pair sequences, two that recognize degenerate six base-pair sequences, and one that recognizes a degenerate five base-pair sequence. The sample of 47 black howler monkeys revealed two mtDNA haplotypes based on the presence or absence of a HINDIII restriction site. In total, 46 individuals exhibited the common profile (PIG1) and a single individual exhibited the variant profile characterized by an extra HINDIII restriction site (PIG2). No mtDNA variability was observed for the remaining 12 enzymes in *A. pigra*.

## DISCUSSION

Levels of allozyme variation in the Bermudian Landing population of Belizean black howler monkeys are low relative to that found in other primates (Table 4). Mean heterozygosity for 36 allozyme loci was 0.021. Other primates and mammals in general tend to have much higher levels of heterozygosity. Pope (1992) found high levels of variation in Venezuelan populations of the red howler *A. seniculus*. Ten out of 29 gene loci were polymorphic ( $P = 32.7$ ), and heterozygosity ranged from 0.057 to 0.135, with a mean  $H$  of 0.099. High levels of heterozygosity are also found in Asian populations of *M. senica*, *M. fuscata*, *M. fascicularis*, and *M. mulatta*, and African populations of *P. hemadryas*, *P. anubis* and *C. aethiops* (Table

4). Low levels of allozyme variation have been reported for some island populations of primates. An island population of the Japanese macaque, *M. fuscata yakui*, had no detectable heterozygosity at 29 putative gene loci, but mainland populations of this species also had relatively low heterozygosity, mean  $H = 0.013$  (Nozawa et al., 1977, 1982). Island populations of the rhesus macaque, *M. mulatta*, also had low heterozygosity relative to mainland populations (Melnick, 1988). Founder events, severe bottlenecks, and continuous genetic drift in small island populations are likely to result in reduced variability relative to mainland populations (Soulé, 1973).

Relative to other primates, the mitochondrial DNA diversity of the Bermudian Landing black howler population also was very low (the percent sequence divergence  $P = 0.2$ ). In a review of mtDNA surveys of eight species of *Macaca* and hominoids, Melnick and Hoelzer (1993) report  $P$  values of 0.9–6.0. Although our estimate of  $P$  is low compared to estimates from other primates, the results of this comparison is problematic. Sequence divergence is expected to increase with geographic isolation and many of the studies reported by Melnick and Hoelzer (1993) were conducted over a broader geographic scale than our study of black howlers. However, even in a study of social groups of toque monkeys (*M. sinica*) that regularly exchange males, there was substantially more sequence divergence

( $P = 3.1$ ) than was observed in black howlers (Hoelzer et al., 1994).

More interindividual differences were noted at the allozyme loci than for the mtDNA, but it is difficult to compare relative levels of genetic diversity because these genetic systems differ in several fundamental ways. Unlinked allozyme loci represent independent genetic markers while mtDNA restriction sites are linked because the molecule is transmitted as a single locus without recombination. Additionally, allozyme loci generally exhibit Mendelian inheritance, but mtDNA generally exhibits maternal transmission. Because of these differences, the effective population size of mtDNA is smaller than the effective population size for nuclear genes such as allozymes (Wilson et al., 1985; Avise et al., 1988). If the population of Black Howlers at Bermudan Landing has been severely reduced in the past, the relatively small effective size of mtDNA means that diversity at this locus would be lost more rapidly than allozyme diversity. Although the almost complete absence of mtDNA diversity in *A. pigra*, taken along with the low levels of allozyme variation, is consistent with the hypothesis that the Bermudian Landing population was drawn through one or more recent severe bottlenecks, we cannot exclude the possibility that low genetic diversity is characteristic of black howler monkey populations, in general. Samples of *A. pigra* from other localities would be necessary to refute the latter hypothesis. A general finding of naturally low levels of variability in other populations would diminish the need for management programs aimed at elevating levels of genetic diversity in locally threatened populations, such as the Bermudian Landing black howlers.

Analysis of population structure with Wright's  $F$ -statistics suggests there is no differentiation among black howler troops. For one of the two loci, there was also a deficiency of heterozygotes. These observations differ dramatically from those made for troops of red howlers (*A. seniculus*) which exhibit excess heterozygosity within populations and significant genetic differentiation among troops (Pope, 1992). Excess heterozygotes within troops with a lack of intertroop

differentiation has also been observed in other primates (Melnick et al., 1984).

In red howler monkeys, females typically remain in their natal troops or occasionally disperse to establish new troops, while dispersing males often immigrate to and reproduce in troops in which they were not born (Pope, 1992). Under this system of social structure, excess heterozygosity within groups and significant differentiation among troops, is the theoretical expectation (Chesser, 1991a; Pope, 1992). It is interesting to speculate why black howlers deviate from these expectations. Past demographic bottlenecks would reduce genetic differentiation among troops (Chesser, 1991a); however, they do not explain the heterozygote deficiencies occurring in most troops. One potential explanation for the heterozygote deficiencies and lack of spatial structure may simply result from our inability to estimate Wright's  $F$ -statistics precisely due to a small number of polymorphic loci in the black howler troops. Interlocus variation in estimates of  $F$ -statistics are often the major source of the variation in these estimates (Nei, 1978). With only two polymorphic loci, it is possible that our measurements of  $F_{IS}$ ,  $F_{IT}$ , and  $F_{ST}$  poorly estimate the true values of these parameters.

Deviation of the  $F$ -statistics from the expectations derived by Chesser (1991a,b) may also result from the demographic bottlenecks the Bermuda landing population has experienced. These exceptions assume populations are at a demographic equilibrium with nonoverlapping generations. This assumption clearly does not apply to the black howler troops we examined, given the recent history of population declines in the CBS. In the most recent documented bottleneck, howler troops were thought to be drastically reduced by a hurricane in 1978 (Bolin, 1981). Following the hurricane, most of the howler troops were comprised of only a single pair of adults. This unusual social structure was assumed to be related to the merging of remnants of troops devastated by the hurricane (Bolin, 1981), suggesting the demographic and genetic structure of the population was disturbed as recently as 1978. Simulations of Chesser (1991a) suggest that a population would reach equilibrial levels

of heterozygosity excess within troops, as well as among troop differentiation, within six generations. Although we have insufficient data to precisely quantify generation length in black howlers, marked individuals do not reproduce until 4–5 years of age, and one female was still reproductively active at 20 years of age. We collected our samples 12 years after the population was greatly reduced by the hurricane. Thus, it is likely that sufficient time has occurred for the population to reach equilibrium conditions with regard to Wright's F-statistics.

Another possibility for the differences between the genetic structure of black howlers and that of red howlers (Pope, 1992) may result from differences in dispersal patterns of these two species. Pope (1992) found that essentially no female successfully immigrated to an established troop after dispersing from its natal troop. We are just beginning to accumulate data on dispersal of marked individuals in the Bermudian Landing population of black howlers (Horwich and James, in prep.), but it appears that some females successfully move among troops (Horwich, 1983a). Between 1990 and 1995, three females successfully immigrated into the 10 troops. The full extent and ramification of both male and female migration will be resolved as more data are collected on the marked individuals. Based on models presented by Chesser (1991a,b), even a small amount of female migration would tend to homogenize gene frequencies among troops. It would have been interesting to study patterns of mtDNA among troops because as a maternally inherited marker, it might have provided additional data on the degree of female philopatry. Unfortunately, the low diversity of mtDNA made this molecule uninformative with regard to the study of population structure.

#### ACKNOWLEDGMENTS

We thank R.H. Horwich, the Community Baboon Sanctuary, the Belizean Ministry of Agriculture, Belize Fisheries, and the Belize Audubon Society for help in the collection and storage of tissue samples. Special thanks also go to J. Brown, K.E. Glander, W. Russell, M. Weinstein, C. Young, and F. Young for field assistance. Also, we thank the past

and present crew at CTAG, Rutgers University for their physical and mental support.

#### LITERATURE CITED

- Avise JC, Ball RM, and Arnold J (1988) Current versus historical population sizes in vertebrate species with high gene flow: A comparison based on mitochondrial DNA lineages and inbreeding theory for neutral mutations. *Mol. Biol. Evol.* 5:331–344.
- Baldwin LA (1976) Vocalizations of howler monkeys (*Alouatta palliata*) in Southwestern Panama. *Folia Primatol.* 26:81–108.
- Bolin I (1981) Male parental behavior in black howler monkeys (*Alouatta palliata*) in Belize and Guatemala. *Primates* 22:349–360.
- Bush GL, Case SM, Wilson AC, and Patton JL (1977) Rapid speciation and chromosomal evolution in mammals. *Proc. Natl. Acad. Sci. USA* 74:3942–3946.
- Chakraborty R, and Nei M (1977) Bottleneck effects on average heterozygosity and genetic distance with the stepwise mutation model. *Evolution* 31:347–356.
- Chesser RK (1991a) Gene diversity and female philopatry. *Genetics* 127:437–447.
- Chesser RK (1991b) Influence of gene flow and breeding tactics on gene diversity within populations. *Genetics* 129:573–583.
- Cohen BJ, and Bree MM (1978) Chemical and physical restraint of nonhuman primates. *J. Med. Primatol.* 7:193–201.
- Glander KE (1980) Reproduction and population growth in free-ranging mantled howling monkeys. *Am. J. Phys. Anthropol.* 53:25–36.
- Glander KE (1991) Capture techniques for three species of monkeys in Costa Rica. *Folia Primat.* 57:70–82.
- Gray JE (1845) On the howling monkeys. *Annals and Magazine of Natural History* 16:217–221.
- Harris H, and Hopkinson DA (1978) *Handbook of Enzyme Electrophoresis in Human Genetics*. Amsterdam: Elsevier.
- Hartshorn GS (1984) Belize Country Environmental Profile a Field Study. Belize City: Nicolait.
- Hayasaka KT (1987) Population genetic study of Japanese macaques. *Macaca fuscata*, in the shimokita a1 troop, with special reference to genetic variability and relationships to Japanese macaques in other troops. *Primates* 28:507–516.
- Hoelzer GA, Dittus WPJ, Ashley MV, and Melnick DJ (1994) The local distribution of highly divergent mitochondrial DNA haplotypes in toque macaques *Macaca sinica* at Polonnaruwa, Sri Lanka. *Molecular Ecology* 3:451–458.
- Horwich RH (1983a) Breeding behaviors in the black howler monkey (*Alouatta pigra*) of Belize. *Primates* 24:222–230.
- Horwich RH (1983b) Species status of the black howler monkey, *Alouatta pigra*, of Belize. *Primates* 24:288–289.
- Horwich RH, and Lyon J (1990) *A Belizean Rain Forest The Community Baboon Sanctuary*. Gay Mills: Oran-utan press.
- Ishimoto G (1973) Blood protein variation in Asian macaques III: Characteristics of the macaque blood polymorphism. *J. Anthropol. Soc. Nippon* 81:1–13.
- Kawamoto Y, Ishcok TM, and Supriatna J (1984) Genetic variations within and between troops of the crab-eating macaque (*Macaca fascicularis*) Sumatra, Java, Bali, Iombok, and Sumbawa, Indonesia. *Primates* 25:131–159.
- Lansman RA, Shade RO, Shapira JF, and Avise JC (1981) The use of restriction endonucleases to measure mitochondrial DNA sequence relatedness in natu-

- ral populations. III. Techniques and potential applications. *J. Mol. Evol.* 17:214–226.
- Lawrence B (1933) Howler monkeys of the *palliata* group. *Bull. Mus. Comp. Zool.* 75:313–354.
- Leberg PL (1992) Effects of population bottlenecks on genetic diversity as measured by allozyme electrophoresis. *Evolution* 46:477–494.
- Maruyama T, and Fuerst PA (1985a) Population bottlenecks and nonequilibrium models in population genetics. III. Genic homozygosity in populations which experience periodic bottlenecks. *Genetics* 111:691–703.
- Maruyama T, and Fuerst PA (1985b) Population bottlenecks and nonequilibrium models in population genetics. II. Number of alleles in a small population that was formed by a recent bottleneck. *Genetics* 111:675–689.
- Melnick DJ (1987) The genetic consequences of primate social organization: A review of macaques, baboons, and vervet monkeys. *Genetica* 73:117–135.
- Melnick DJ (1988) The genetic structure of a primate species: Rhesus macaques and other cercopithecine monkeys. *Internat J. Primatol.* 9:195–231.
- Melnick DJ, and Hoelzer GA (1993) What is mtDNA good for in the study of primate evolution. *Evolutionary Anthropol.* 9:195–231.
- Melnick DJ, Jolly CJ, and Kidd KK (1984) The genetics of a wild population of rhesus monkeys (*Macaca mulatta*). I. Genetic variability within and between social groups. *Am. J. Phys. Anthropol.* 63:341–360.
- Murphy RW, Sites JW Jr., Buth DG, and Haufler CH (1990) Proteins I: Isozyme Electrophoresis. In D. Hillis and D. Moritz (eds.): *Molecular Systematics*. Sunderland: Sinauer, pp. 45–126.
- Nei M (1977) F-statistics and analysis of genetic diversity in subdivided populations. *Ann. Hum. Genet.* 41:225–233.
- Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89:583–590.
- Nei M (1987) *Molecular Evolutionary Genetics*. New York: Columbia University Press.
- Nei M and Li CC (1979) Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA.* 76:5259–5273.
- Nei M, Maruyama T, and Chakraborty R (1975) The bottleneck effect and genetic variability in populations. *Evolution* 29:1–10.
- Nozawa K, Shotake T, Kawamoto Y, and Tanabe Y (1982) Population genetics of Japanese monkeys: II. Blood protein polymorphisms and population structure. *Primates* 23:252–271.
- Nozawa K, Shotake T, Ohkura Y, and Tanabe Y (1977) Genetic variation within and between species of Asian macaques. *Jap. J. Genetics* 52:15–30.
- Pope TR (1992) The influence of dispersal patterns and mating system on genetic differentiation within and between populations of the red howler monkey (*Alouatta seniculus*). *Evolution* 46:1112–1128.
- Scott NJ Jr, Scott AF, and Malmgren LA (1976) Capturing and marking howler monkeys for field behavioral studies. *Primates* 17:527–533.
- Shotake T (1981) Population genetical study of natural hybridization between *Papio anubis* and *P. hamadryas*. *Primates* 22:285–308.
- Shotake T, and Nozawa K (1984) Blood protein variations in baboons. II. Genetic variability within and among herds of gelada baboons in the Central Ethiopian plateau. *J. Hum. Evol.* 13:265–274.
- Shotake T, and Santiapillai C (1982) Blood protein polymorphism in the troops of the toque macaques, *Macaca sinica*, in Sri Lanka. *Kyoto University Overseas Research Report of Studies on Asian Non-human Primates* 2:79–95.
- Sirkkomaa S (1983) Calculations on the decrease of genetic variation due to the founder effect. *Hereditas* 99:11–20.
- Smouse PE, and Neel JV (1981) The impact of random and lineal fission on the genetic divergence of small human groups: A case study among the Yanomama. *Genetics* 98:179–197.
- Soulé M (1973) The epistasis cycle: A theory of marginal populations. *Ann. Rev. Ecol. Syst.* 4:165–187.
- Southern EM (1975) Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98:503–517.
- Swofford DL, and Selander RK (1981) BIOSYS-1: A Fortran program for the comprehensive analysis of electrophoretic data in population genetics and systematics. *J. Heredity* 72:281–283.
- Turner TR (1981) Blood protein variation in a population of Ethiopian vervet monkeys (*Cercopithecus aethiops aethiops*). *Am. J. Phys. Anthropol.* 55:225–232.
- Weir BS, and Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. *Evolution* 38:1358–1370.
- Wilson AC, Cann RL, Carr SM, George M, Gyllenstein UB, Helm-Bychowski KM, Higuchi RG, Palumbi SR, Prager EM, Sage RD, and Stoneking M (1985) Mitochondrial DNA and two perspectives on evolutionary genetics. *Biol. J. Linnean Soc. London* 26:375–400.
- Workman PL, and Niswander (1970) Population studies on southwestern Indian tribes. II. Local genetic differentiation in the Papago. *Am. J. Hum. Genet.* 22:724–749.
- Wright S (1965) The interpretation of population structure by f-statistics with special regard to systems of mating. *Evolution* 19:395–420.